

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/47, C07D 491/22	A1	(11) International Publication Number: WO 90/03169 (43) International Publication Date: 5 April 1990 (05.04.90)
(21) International Application Number: PCT/US89/04176 (22) International Filing Date: 28 September 1989 (28.09.89) (30) Priority data: 250,094 28 September 1988 (28.09.88) US (71) Applicant: RESEARCH TRIANGLE INSTITUTE [US/ US]; P.O. Box 12194, Research Triangle Park, NC 27709-2194 (US). (72) Inventors: WALL, Monroe, E. ; 1826 North Lakeshore Drive, Chapel Hill, NC 27514 (US). WANI, Mansukh, C. ; 2801 Legion Avenue, Durham, NC 27707 (US). NI- CHOLAS, Allan, W. ; 1001 Millbrook Road, Raleigh, NC 27609 (US). MANIKUMAR, Govindarajan ; 2563 New Hope Church Road, Raleigh, NC 27604 (US).	(74) Agents: OBLON, Norman, F. et al.; Oblon, Spivak, McClelland, Maier & Neustadt, Fourth Floor, 1755 South Jefferson Davis Highway, Arlington, VA 22202 (US). (81) Designated States: AT (European patent), AU, BE (Euro- pean patent), CH (European patent), DE (European pa- tent), DK, FR (European patent), GB (European pa- tent), IT (European patent), JP, KR, LU (European pa- tent), NL (European patent), NO, SE (European patent). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>	
(54) Title: SYNTHESIS OF CAMPTOTHECIN AND ANALOGS THEREOF (57) Abstract A method for synthesizing camptothecin and camptothecin analogs using a novel hydroxyl-containing tricyclic intermedi- ate and the camptothecin analogs produced by the process. The camptothecin analogs are effective inhibitors of topoisomerase I and show anti-leukemic and anti-tumor activity.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LJ	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

-1-

DescriptionSynthesis of Camptothecin and Analogs ThereofTechnical Field

The invention relates to camptothecin and analogs
5 thereof which show life prologation effects in various
leukemia systems such as P-388 and L-1210; inhibition of
animal tumors such as B-16 melanoma and are potent
inhibitors of topoisomerases I and II. It also relates
to a method of synthesizing the same by means of a novel
10 hydroxyl-containing tricyclic intermediate.

Background Art

Camptothecin is a pentacyclic alkaloid initially
isolated from the wood and bark of Camptotheca acuminata
by Wall et al (M.E. Wall, M.C. Wani, C.E. Cook, K.H.
15 Palmer, A.T. McPhail, and G.A. Sim, J. Am. Chem. Soc.,
94, 388 (1966).

Camptothecin is highly biologically active and
displays strong inhibitory activity toward the
biosynthesis of nucleic acids. Additionally,
20 camptothecin exhibits potent anti-tumor activity against
experimentally transplanted carcinoma such as leukemia L-
1210 in mice or Walker 256 tumor in rats.


Several methods for the synthesis of camptothecin
and camptothecin analogs are known. These synthetic
25 methods include (i) methods in which naturally occurring
camptothecin is synthetically modified to produce a
number of analogs and (ii) totally synthetic methods.

U.S. Patents 4,604,463; 4,545,880; and 4,473,692 as
well as European Patent Application 0074256 are examples
30 of the former type of synthetic strategy. Additional
examples of this strategy can be found in Japanese
Patents 84/46,284; 84/51,287; and 82/116,015. These

methods require naturally occurring camptothecin which is difficult to isolate and hence these methods are not suitable for the production of large quantities of camptothecin or analogs.

5 Examples of a variety of totally synthetic routes to camptothecin and camptothecin analogs can be found in the following references: Sci. Sin. (Engl. Ed), 21(1), 87-98 (1978); Fitoterapia, 45(3), 87-101 (1974); Yakugaku Zasshi, 92(6), 743-6 (1972); J. Org. Chem., 40(14), 2140-1
10 (1975); Hua Hsueh Hsueh Pao, 39(2), 171-8 (1981); J. Chem. Soc., Perkin Trans 1, (5), 1563-8 (1981); Heterocycles, 14(7), 951-3 (1980); J. Amer. Chem. Soc., 94(10), 3631-2 (1972); J. Chem. Soc. D, (7), 404 (1970) and U.S. Patent 4,031,098.

15 Wani et al, J. Med. Chem., 23, 554 (1980) discloses a synthesis of camptothecin and camptothecin analogs which involves the reaction of a tricyclic compound with a suitably substituted ortho-aminoaldehyde to yield desoxycamptothecin as shown in Equation 1 below.



Desoxycamptothecin is then treated with oxygen to give camptothecin analogs. A major disadvantage of this procedure is the insolubility of desoxycamptothecin and
25 its analogs, requiring large solvent volumes in the final step. A poor yield of the oxygenation product results under these conditions.

There exists a need, therefore, for a high-yield,

efficient synthesis of camptothecin and camptothecin analogs which does not require prior isolation of naturally occurring camptothecin.

A need also exists for a method of synthesizing
5 camptothecin and camptothecin analogs which does not suffer from insolubility problems of intermediate compounds and the resulting low yields.

A further need exists for new camptothecin analogs which can be synthesized in an efficient, high-yield
10 manner and which show good biological activity.

Disclosure of the Invention

Accordingly, one object of the present invention is to provide a method of synthesizing camptothecin and camptothecin analogs in high yield in a totally synthetic
15 process.

Another object of the present invention is to provide a process for synthesizing camptothecin and camptothecin analogs which does not suffer from problems associated with the insolubility of intermediate
20 compounds.


A further object of the invention is to provide a process for the preparation of camptothecin and camptothecin analogs which can be easily modified to produce a variety of analog structures.

25 Still a further object of the present invention is to provide camptothecin analogs which show good antitumor activity and other desirable biological activities.


These objects and other objects of the present invention which will become apparent from the following
30 specification have been achieved by the present method for the synthesis of camptothecin and camptothecin analogs, which includes the steps of:

cyclizing a compound of the formula shown below,
wherein X is an organic group which is converted to a


carbonyl group when treated with an acid,



to form a lactone having the formula



- 5 deprotecting said lactone to form a hydroxyl-containing tricyclic compound having the formula shown below, and



- reacting said hydroxyl-containing tricyclic compound
10 with a substituted ortho-amino compound of the formula



- wherein $n = 1-2$ and wherein each R is selected from the group consisting of cyano, methylenedioxy, formyl, hydroxy, C_{1-8} alkoxy, nitro, amino, chloro, bromo, iodo, 15 fluoro, C_{1-8} alkyl, trifluoromethyl, aminomethyl, azido, amido and hydrazino groups; R^2 is H_1 or C_{1-8} alkyl; and R^3 is the side-chain of any of the twenty naturally occurring amino acids.


BRIEF DESCRIPTION OF THE DRAWING

- 20 Figures 1a and 1b illustrates the synthesis of the hydroxyl-containing tricyclic compound 11, according to the invention.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The camptothecin and camptothecin analogs produced by the process of the present invention are racemic, and therefore contain both the (R)-20-hydroxy and (S)-20-hydroxy camptothecin compounds. Naturally occurring camptothecin belongs to the 20(S) series of compounds. Therefore, the compounds produced by the process of the present invention contain a mixture of the natural and non-naturally occurring compounds.

The camptothecin analogs of the present invention have the basic camptothecin structural framework shown below in which the A ring is substituted.



5 Substituents within the scope of the present invention include hydroxy, nitro, amino, chloro, bromo, iodo, fluoro, C₁₋₈ alkyl, C₁₋₈ alkoxy, trifluoromethyl, aminomethyl, amido, hydrazino, azido, formyl, and cyano groups as well as groups comprising amino acids and/or
 10 peptides bonded to the aromatic ring via the amino-nitrogen atom or via an amide linkage which contains the carbonyl group of the amino acid, i.e., an amino acid amido group. Amides prepared from cyclic anhydrides may also be prepared. Preferred alkyl groups include methyl,
 15 ethyl, propyl, butyl, isopropyl, isobutyl and sec-butyl groups. Preferred alkoxy groups include methoxy, ethoxy, propoxy and isopropoxy groups.

The preferred amino acid groups are the 20 naturally occurring amino acids having an (L) configuration. These
 20 amino acids are well known to those skilled in the art.

Water-soluble camptothecin analogs which can be prepared by the process of the present invention include analogs in which R_n is an amide linkage formed by reacting an amino-camptothecin analog with the free
 25 carboxylic acid group of an amino acid, peptide or carboxylic acid derivative thereof. These amide derivatives may be present as the free amines, i.e., the α-amino group or as any of the well known acid addition salts, such as, for example, hydrochloride,
 30 gluconate, phosphate or hydrobromide addition salts. Additional water soluble analogs can be prepared by reacting an amino-camptothecin analog with a cyclic carboxylic acid anhydride to give a carboxylic acid amide group. This reaction results in an amide substituent on
 35 the camptothecin structural framework and a free

carboxylic acid group which may be present as the free acid or as a salt, e.g., alkali metal, or alkaline-earth metal salt, or as an organic cation, e.g., ammonium salts. Preferred anhydrides are the C₄₋₁₀ saturated and 5 unsaturated acid anhydrides. Alternatively, the above water soluble derivatives can also be prepared by employing the corresponding ester-acid halide in place of the anhydride. Water soluble urea and urethane analogs can also be prepared by reacting an amino-camptothecin 10 analog with phosgene followed by an appropriate diamine, e.g. n-alkylpiperazine or an appropriate tertiary-amino alcohol, e.g. n-dialkyl-aminoethanol respectively. These water soluble derivatives are particularly important since they may be therapeutically administered 15 in aqueous pharmaceutical compositions.

Additionally, water soluble dialkylamino ether analogs of camptothecin can be prepared by reacting an appropriate tertiary-aminoalkyl halide, e.g. dialkylaminoethyl halide with an hydroxy-camptothecin 20 analog followed by salt formation as described above.


Additionally, two substituents on the A ring may be joined together to form a bifunctional substituent such as the methylenedioxy group. Methylenedioxy substituents may be bonded to any two consecutive positions in the A 25 ring, for example, the 9,10; 10,11 or 11,12 positions.

Preferred substituents include the hydroxy, amino, cyano, methylenedioxy, 9 or 10-glycinamido, 9- or 10-succinamido, 9- or 1-(4-methylpiperazine) carbonylamino, 9- or 10-(N,N-diethylaminoethoxy) carbonylamino, and 9- 30 diethylaminoethoxy substituents. A particularly preferred substituent is the methylenedioxy group.

Particularly preferred compounds within the scope of the invention include 11-methoxy-20(RS)-camptothecin, 11-hydroxy-20(RS)-camptothecin, 10-hydroxy-20(RS)- 35 camptothecin, 9-methoxy-20(RS)-camptothecin, 9-hydroxy-20(RS)-camptothecin, 10-nitro-20(RS)-camptothecin, 10-amino-20(RS)-camptothecin, 9-nitro-20(RS)-camptothecin,

9-amino-20(RS)-camptothecin, 11-nitro-20(RS)-
camptothecin, 11-amino-20(RS)-camptothecin, 10,11-
dihydroxy-20(RS)-camptothecin, 10-chloro-20(RS)-
camptothecin, 10-methyl-20(RS)-camptothecin, 11-formyl-
5 20(RS)-camptothecin and 11-cyano-20(RS)-camptothecin,
10,11-methylenedioxy-20(RS)-camptothecin, 9-glycinamido-
20(RS)-camptothecin and 9-succinamido-20(RS)-
camptothecin, 9- or 10-(4-methylpiperazine)
carbonylamino-20(RS)-camptothecin, 9- or 10-(N,N-
10 diethylaminoethoxy) carbonylamino-20(RS), camptothecin,
and 9-diethylaminoethoxy-20(RS)-camptothecin.


Also included within the scope of the present
invention are compounds in which the A ring of the
camptothecin structure is modified to contain a hetero
15 atom. The modified structures can have an A ring which
contains 5 or 6 atoms and the hetero atom may be a
nitrogen, sulfur or oxygen atom. These compounds may be
represented by the general structure shown below in which
the A ring is an aromatic 5 or 6 membered ring containing
20 the hetero atom X.



Preferred compounds having a modified A ring structure include compounds in which the A ring is a 6 membered nitrogen-containing aromatic ring and compounds in which the A ring is a 5 membered sulfur-containing aromatic ring. Particularly preferred compounds are 10-aza-20(RS)-camptothecin and A-nor-9-thia-20(RS)-camptothecin.

The camptothecin analogs noted above may be synthesized according to the method of the present invention by reacting a tricyclic compound containing a 20-hydroxyl group with an appropriately substituted ortho-amino aromatic aldehyde or ketone. Camptothecin analogs having an alkyl substituent on C₇ are produced when the appropriate ortho-amino ketone is used.

An important step in the method of the present invention is the synthesis of the hydroxyl-containing tricyclic compound having the formula I shown below and in which R is a hydroxyl group.



(I)

A synthetic method previously developed by the present inventors (J. Med. Chem., 23, 554 (1980)) utilized a related but structurally different tricyclic compound (formula I, R = H). In that method, the tricyclic compound was reacted with a suitable ortho-aminoaldehyde under alkaline or acidic conditions to yield a desoxycamptothecin. The desoxycamptothecin was then reacted with oxygen to give camptothecin analogs in which R is OH. A major disadvantage of this procedure is the insolubility of the desoxycamptothecin and its analogs, requiring large solvent volumes in the final

step and giving poor yields of the oxygenation product.


In contrast, the method of the present invention synthesizes the key tricyclic intermediate (11) according to Figure 1. The synthesis of compounds 1-9 was disclosed in Wani et al, J. Med. Chem., 23, 554 (1980). In further contrast to the previous synthesis, the present method introduces the 20-hydroxyl group earlier in the synthetic sequence and then forms the lactone ring to give compound 10. After deprotection of the carbonyl group, the key hydroxyl-containing tricyclic compound 11 is obtained.

The protection of the carbonyl group in compound 3 can be performed using any appropriate organic protecting group which can be removed or converted into a carbonyl group upon treatment with acid. The carbonyl group is thereby "deprotected". These protecting groups are well known to those familiar with synthetic chemistry, and include acetals, ketals, thioacetals, thioketals, etc. Preferred protecting groups have 2-6 carbon atoms. An especially preferred protecting group is $-OCH_2CH_2O-$.

As a consequence of prior introduction of the hydroxyl group into the tricyclic compound 11, the desired pentacyclic analogs are produced in one step by reaction with the appropriate ortho-amino carbonyl compounds. Both compound 11 and the corresponding ketonic synthons are very soluble in organic solvents whereas the pentacyclic product is insoluble. Hence, the oxygenation step, i.e, the introduction of the hydroxyl group, is conveniently carried out at the tricyclic stage rather than on the insoluble pentacyclic desoxy analogs.

Tricyclic compound 11 is then reacted with a suitably substituted ortho-amino aldehyde or ketone to give a camptothecin analog. Substituted ortho-amino aldehydes and ketones within the scope of the present invention include ortho-amino aldehydes and ketones having at least one additional substituent on the aromatic ring. This substituent may be at one or more of

the positions equivalent to the 9, 10, 11 or 12 positions of the A ring of the final camptothecin structure as shown below.




5 Preferred substituted ortho-amino aldehydes and ketones have substituents in one or more of the equivalent 9, 10, or 11 positions.

The substituents on the substituted ortho-amino-aldehyde or ketone include hydroxy, nitro, amino, C₁-8
 10 alkyl, chloro, bromo, iodo, fluoro, methylenedioxy (-O-CH₂-O-), C₁-8 alkoxy, trifluoromethyl, aminomethyl, amido, hydrazino, azido, formyl, and cyano groups as well as groups comprising amino acids bonded to the aromatic ring through the amino-nitrogen atom. Preferred examples
 15 include the hydroxy, amino, cyano and methylenedioxy substituents. A particularly preferred substituent is the methylenedioxy group.

When an ortho-amino ketone is reacted with tricyclic compound 11, a camptothecin analog having an alkyl
 20 substituent at C₇ is produced. Preferred ortho-amino ketones are those in which R² is an alkyl group having 1-8 carbon atoms. Especially preferred ortho-amino ketones are ortho-aminoacetophenone and ortho-aminopropiophenone.

The ortho-amino aldehydes and ketones may be
 25 substituted by a group having the formula



wherein R³ is a side-chain of one of the twenty naturally occurring amino acids. The amino acid substituent is bonded to the aromatic ring via the nitrogen atom and may
 30 be bonded to any position on the aromatic ring equivalent to the 9, 10, 11 or 12 positions of the A ring of the final camptothecin structure.

The ortho-amino aldehydes and ketones may be in the free carbonyl form or in a form in which the carbonyl of the aldehyde or ketone is protected by a standard protecting group. These protecting groups are well known to those skilled in the art. Ortho-amino aldehydes and ketones in the free carbonyl form and in the protected carbonyl form are considered within the scope of the present invention and are suitable for use in the present method.

10 The reaction in which the hydroxyl group is introduced into the tricyclic intermediate compound, i.e., the cyclizing step, can be effected by any suitable reaction which will introduce the hydroxyl group at the appropriate position of compound 9 without causing
15 significant side reactions such as degradation of compound 9 itself.

The reaction is preferably conducted in the presence of a basic catalyst. Suitable basic catalysts include both inorganic and organic bases. Preferred inorganic
20 bases include, for example, sodium and potassium carbonate and sodium and potassium bicarbonate. Preferred organic bases include hindered bases such as triethylamine and diisopropylamine. A particularly preferred basic catalyst is potassium carbonate.

25 The reaction in which the hydroxyl group is introduced can be performed in the presence of any polar or non-polar solvent in which the reactants are suitably soluble to react. Preferred are polar organic solvents such as methanol, ethanol, propanol, butanol and
30 dimethylformamide. Ether solvents, including crown ethers may also be used.

The oxygen of the hydroxyl group is generally derived from molecular oxygen which is bubbled through the reaction solution. Although the use of oxygen is
35 preferred, other sources of oxygen, such as air, may also be used. Other oxidizing agents such as hydrogen peroxide, lead tetraacetate and selenium dioxide may also

be employed.

This reaction is preferably conducted at room temperature although the specific reaction temperature will be dependent on the specific reaction conditions and 5 reactants used.

The deprotection of the carbonyl group in compound 10 is accomplished by treatment with acid. Suitable acids include mineral acids such as HCl, H₂SO₄, HNO₃, and H₃PO₄, as well as organic acids such as alkanolic acids 10 having 1-10 carbon atoms, preferably acetic acid, and C₁₋₁₂ arylsulfonic acids, especially p-toluenesulfonic acid. The deprotection of a carbonyl group in this manner is well known to those skilled in the art.

The tricyclic compound 11 is then reacted with a 15 substituted ortho-amino aldehyde or ketone in the presence of an acid or base catalyst. The base catalyst is preferably any of the base catalysts noted above in cyclizing compound 9 to form compound 10, i.e., for the introduction of the hydroxyl group into tricyclic 20 compound 11. The acid catalyst is preferably a mineral acid such as for example HCl, H₂SO₄, HNO₃, and H₃PO₄, or organic acids such as C₁₋₈ alkanolic acids and C₁₋₁₂ arylsulfonic acids, especially p-toluenesulfonic acid.

The reaction of compound 11 with an appropriate 25 ortho-amino compound may be carried out neat or in the presence of a polar or non-polar solvent. Preferred polar solvents are the C₁₋₆ alcohols, ethers and dimethylformamide. Preferred non-polar solvents are branched or straight chained alkyl hydrocarbons having 4- 30 10 carbon atoms and aromatic hydrocarbons having 6-20 carbon atoms. An especially preferred solvent is toluene.

The reaction of the hydroxyl-containing tricyclic compound with the optionally substituted ortho-amino 35 compound is generally conducted with heating at reflux. Reaction times will vary depending on the particular

reactants but are generally in the range from about 10 minutes to 24 hours. Preferred reaction times are in the range of 2-10 hours.

The camptothecin analogs of the present invention
5 have excellent biological activity. As used herein, "biological activity" refers to the ability of the camptothecin analogs to inhibit topoisomerase enzymes, in particular topoisomerase I, and their ability to exert anti-leukemic activity. Anti-leukemic activity may be
10 determined by the ability of the respective compounds to inhibit L-1210 mouse leukemia cells. Although anti-leukemic activity is demonstrated here by the activity of the particular compounds against L-1210 mouse leukemia cells, other known anti-leukemic and anti-tumor in vitro
15 and in vivo models may be used as well to determine anti-leukemic activity.

The mouse anti-leukemic activity of the various ring A oxygenated camptothecin analog is shown in Table I. Similar data for nitrogen analogs and for ring A modified
20 analogs are shown in Tables II and III, respectively. In most cases camptothecin or an analog with well-defined activity was also assayed at the same time as a positive control, and the data are shown in the table footnotes. In this manner the relative anti-leukemic activity of the
25 various compounds can be compared. The biological activity of additional camptothecin analogs is described in J. Med. Chem., 23, pages 554-560 (1980) incorporated herein by reference.

The ability of camptothecin to inhibit topoisomerase
30 I has been shown. See J. Biol. Chem., 260, 14873-73 (1985) incorporated herein by reference.

Table I. Comparative Activities and Potencies of Ring A Oxygenated Camptothecin Analogues in Mouse Leukemia Assays^{a,b}

Camptothecin derivative	max % T/C (dose, mg/kg)	no. cures out of 6	K _E ^c at max % T/C	active dose range, mg/kg	toxic dose, mg/kg
10-OH-20(S) ^{a,d}	297 (3.1)	0		0.4 ^e -3.1	6.25
10-OMe-20(S) ^{a,d}	167 (1.6)	0		0.4 ^e -1.6	3.1
11-OH-20(RS) ^{b,f}	357 (60.0)	3	J 5.68	7.5 ^e -60.0 ^g	>60.0
10,11-diOMe-20(RS) ^{b,h}	inactive				>50.0 ^g
10,11-OCH ₂ O-20(RS) ^{b,i}	325 (2.0)	2	J 5.97	2.0 ^e -4.0	>8.0
10-OCH ₂ CO ₂ Na-20(S) ^{a,d}	inactive				
10-Et ₂ N(CH ₂) ₂ O-20(S) ^{a,d}	183 (16.0)	0		2.0 ^e -32.0 ^g	>32.0

^aDenotes testing in P-388 system; treatment schedule Q04DX03; %T/C = survival time of treated/control animals X 100; IP using Klucel emulsifier. ^bDenotes testing in L-1210 system; treatment schedule Q04DX02; %T/C = survival time of treated/control animals X 100; IP using Klucel emulsifier. ^cLog₁₀ of initial tumor cell population minus log₁₀ of tumor cell population at end of treatment. ^d20(S)-Camptothecin and 20(S)-camptothecin sodium were used as reference standards; for 20(S)-camptothecin, %T/C (4.0 mg/kg) = 197; for 40(S)-camptothecin sodium, %T/C (40.0 mg/kg) = 212. ^eLowest dose administered. ^f20(S)-camptothecin and 20(S)-camptothecin sodium were used as reference standards: for 20(S)-camptothecin, %T/C (8.0 mg/kg) = 164; for 20(S)-camptothecin sodium, %T/C (40.0 mg/kg) = 178. ^gHighest dose administered. ^h20(S)-camptothecin sodium was used as a reference standard: %T/C (25.0 mg/kg) = 206. ⁱ20(S)-camptothecin was used as a reference standard: %T/C (5 mg/kg) = 166.

Table II. Comparison of Activities and Potencies of Ring A Nitrogen Substituted and Ring A Nitrogen/Oxygen Disubstituted Analogues in L-1210 Mouse Leukemia Assays^a

Camptothecin derivative	max % T/C (dose, mg/kg)	no. cures out of 6	K _E ^b at max % T/C	active dose range, mg/kg	toxic dose, mg/kg
10-NO ₂ -20(RS) ^c	219 (15.5)	1	J 5.86	7.5 ^d -15.5	31.0
10-NH ₂ -20(RS) ^c	329 (8.0)	3	J 5.86	4.0 ^d -16.0	32.0
10-NHAc-20(RS) ^c	318 (40.0)	1	J 5.86	5.0 ^d -40.0 ^e	>40.0
9-NO ₂ -20(S) ^f	348 (10.0)	5	J 5.86	2.5 ^d -20.0	40.0 ^e
12-NO ₂ -20(S) ^f	151 (40.0)	0	0.34	2.5 ^d -40.0 ^e	>40.0
9-NH ₂ -20(S) ^f	348 (2.5)	4	J 5.86	2.5 ^d -5.0	10.0
12-NH ₂ -20(S) ^f	inactive				
9-NO ₂ -10-OMe-20(S) ^f	160 (40.0)	0	1.03	2.5 ^d -40.0 ^e	>40.0
9-NH ₂ -10-OMe-20(S) ^f	186 (40.0)	0	2.92	2.5 ^d -40.0 ^e	>40.0
9-NO ₂ -10-OH-20(S) ^f	131 (20.0)	0	-1.12	2.5 ^d -40.0 ^e	>40.0
9-NHAc-10-OH-20(S) ^f	220 (40.0)	0	5.50	2.5 ^d -40.0 ^e	>40.0

^aTreatment schedule Q04DX2; % T/C = survival time of treated/control animals X 100; IP using Klucel emulsifier. ^bLog₁₀ of initial tumor cell population minus log₁₀ of tumor cell population at end of treatment. ^c20(S)-Camptothecin (1) and 10-hydroxy-20(S)-camptothecin (2) were used as reference standards: for 1, % T/C (8.0 mg/kg) = 197; for 2, % T/C (24.0 mg/kg) = 230. ^dLowest dose administered. ^eHighest dose administered. ^fCompounds 1 and 2 were used as reference standards: for 1, % T/C (10.0 mg/kg) = 267; for 2, % T/C (20.0 mg/kg) = 348.

Table III. Comparative Activities of Ring A Modified and Homologated Camptothecin Analogues in Mouse Leukemia Assays^{a,b}

Camptothecin derivative	max % T/C (dose, mg/kg)	no. cures out of 6	K _E ^c at max % T/C	active dose range, mg/kg	toxic dose, mg/kg
10-aza-20(RS) ^{a,d}	162 (2.5)	0	1.61	1.25 ^e -2.5	5.0
12-aza-20(RS) ^{b,f}	175 (32.0)	0		8.0-32.0	>32.0
A-nor-9-thia-20(RS) ^{a,d}	193 (25.0)	0	-0.94	3.12 ^e -25.0	50.0 ^g

^a Denotes testing in L-1210 system; treatment schedule Q04DX2; % T/C survival time of treated/control animals X 100; IP using Klucel emulsifier. ^b Denotes testing in P-388 system; treatment schedule Q04DX3; % T/C = survival time of treated/control animals X 100; IP using Klucel emulsifier. ^c Log₁₀ of initial tumor cell population minus log₁₀ of tumor cell population at end of treatment. ^d 20(S)-Camptothecin sodium was used as a reference standard: % T/C (40.0 mg/kg) = 215. ^e Lowest dose administered. ^f 20(S)-Camptothecin was used as a reference standard: % T/C (4.0 mg/kg) = 197. ^g Highest dose administered.

Pharmaceutical compositions containing the novel camptothecin analogs are also within the scope of the present invention. These pharmaceutical compositions may contain any quantity of a camptothecin analog which is
5 effective to inhibit topoisomerase I in vitro or in vivo or exhibit anti-leukemic activity in vivo. Mammals such as humans are treatable with the inventive compositions. Typical in vivo doses within the scope of the invention are from 0.1-60 mg of camptothecin analog
10 per kg of body weight. A particularly preferred range is 1-40 mg/kg.

There may also be included as part of the composition pharmaceutically compatible binding agents, and/or adjuvant materials. The active materials can also
15 be mixed with other active materials which do not impair the desired action and/or supplement the desired action. The active materials according to the present invention can be administered by any route, for example, orally, parenterally, intravenously, intradermally,
20 subcutaneously, or topically, in liquid or solid form.

A preferred mode of administration of the compounds of this invention is oral. Oral compositions will generally include an inert diluent or an edible carrier. They may be enclosed in gelatin capsules or
25 compressed into tablets. For the purpose of oral therapeutic administration, the aforesaid compounds may be incorporated with excipients and used in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gums and the like. These preparations
30 should contain at least 0.1% of active compound but may be varied depending upon the particular form.

The tablets, pills, capsules, troches and the like may contain the following ingredients: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an
35 excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, corn starch and the like; a lubricant such as magnesium stearate or Sterotes;

a glidant such as colloidal silicon dioxide; and a sweetening agent such as sucrose or saccharin or flavoring agent such as p ppermint, methyl salicylate, or orange flavoring may be added. When the dosage unit form is a capsule, it may contain, in addition to material of the above type, a liquid carrier such as a fatty oil. Other dosage unit forms may contain other various materials which modify the physical form of the dosage unit, for example, as coatings. Thus tablets or pills may be coated with sugar, shellac, or other enteric coating agents. A syrup may contain, in addition to the active compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors. Materials used in preparing these various compositions should be pharmaceutically pure and non-toxic in the amounts used.

For the purposes of parenteral therapeutic administration, the active ingredient may be incorporated into a solution or suspension.

The solutions or suspensions may also include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

The dosage values will vary with the specific severity of the disease condition to be alleviated. Good results are achieved when the compounds described herein are administered to a subject requiring such treatment as an effective oral, parenteral or intravenous dose. It is

to be understood that for any particular subject, specific dosage regimens should be adjusted to the individual need and the professional judgment of the person administering or supervising the administration of the aforesaid compound. It is to be further understood that the dosages set forth herein are exemplary only and they do not limit the scope or practice of the invention. The dosages may be administered at once, or may be divided into a number of smaller doses to be administered at varying intervals of time.

Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

15

EXAMPLESSYNTHESIS OF TRICYCLIC COMPOUND 11:

6-Cyano-7-methyl-1,5-dioxo- $\Delta^{6(8)}$ -tetrahydroindolizine (Compound 3).

Ethyl acetopyruvate was prepared from acetone and diethyl oxalate as described in Org. Synthesis, Coll. Vol. 1, 238 (1958). Further reaction with triethylorthoformate and ammonium chloride in ethanol afforded the known enol ether 1. See L. Claisen, Chem. Ber., 40, 3903 (1907).

25 Ethyl(2-ethoxy-4-oxo)-pent-2-enoate(1) (100.01 g, 0.538 mol) was added gradually to a preheated (45°), mechanically stirred mixture of K_2CO_3 (79.04 g, 0.573 mol) and cyanoacetamide (48.46 g, 0.577 mol) in DMF (960 mL). The mixture was kept at 45° for 18 h, whereupon the thick, red slurry was treated dropwise with freshly distilled methyl acrylate (360 mL, 343 g, 3.99 mol). After 72 hr at 45°, the red suspension was filtered, dissolved in 5 liters of water, and acidified to pH 1.5 with concentrated HCl. Crude bicyclic ester 2 35 (127.98 g) was collected by filtration as a pink solid.

Without further treatment, 2 was refluxed in a solution concentrated of HCl (800 ml) and glacial HOAc (800 ml) for 2 hr. Removal of the solvents in vacuo gave the bicyclic pyridone 3 (39.66 g, 39% based on 1).

5 6-Cyano-1,1-(ethylenedioxy)-7-methyl-5-oxo- $\Delta^{6(8)}$ -tetrahydroindolizine (Compound 4).

Compound 3 (10.54 g, 0.056 mol) as a stirred solution in CH_2Cl_2 (500 ml) was treated at room temperature under N_2 with ethylene glycol (6.85 ml, 10 7.63 g, 0.123 mol), and Me_3SiCl (31.30 ml, 26.89 g, 0.247 mol) and left at ambient temperature (20°) for 65 hr. The solution was filtered to remove some black suspended material before washing with 1 M aq NaOH solution. The organic phase was washed with brine, 15 filtered through Celite and evaporated to afford ethylene ketal 4 (10.26 g, 79%) as a pink solid.

6-Cyano-1,1-(ethylenedioxy)-7-[(ethoxycarbonyl)methyl]-5-oxo- $\Delta^{6(8)}$ -tetrahydroindolizine (Compound 5).

The ketal 4 (5.0 g, 0.022 mol) was refluxed in a 20 suspension of KH (11.9 g, 0.068 mol) in toluene (40 mL) for 10 min. Diethyl carbonate (6.79 g, 0.058 mol) and a catalytic amount (0.31 g, 6.7 mmol) of absolute ethanol were added and refluxing continued for 3 hr. The dark solid was crushed and the resulting suspended salt of 5 25 was collected by filtration. The salt was neutralized by the careful addition of cold aqueous HOAc. Water was added and the product extracted into CH_2Cl_2 . Following a wash with brine and drying (Na_2SO_4), evaporation of the CH_2Cl_2 afforded crude 5. Purification by silica gel 30 chromatography (2% MeOH in CHCl_3) and recrystallization (MeOH) gave pure 5 (4.97 g, 76%).

6-Cyano-1,1-(ethylenedioxy)-7-[1'-(ethoxycarbonyl)-propyl]-5-oxo- $\Delta^{6(8)}$ -tetrahydroindolizine (Compound 6).

A stirred solution of the ester 5 (4.01 g,

0.0132 mol) in anhydrous DME (70 mL) at -78°C was treated with potassium tert-butoxide (1.7 g, 15 mmol). After 5 min, EtI (8.24 g, 0.053 mol) was added over a 5 min period. After stirring for 1.5 hr at -78°C, the mixture 5 was left to warm to room temperature overnight. Water was added and the product extracted into CH₂Cl₂. After washing with brine and drying (Na₂SO₄), CH₂Cl₂ was evaporated to give the ester 6 (4.3 g, 98%).

6-(Acetamidomethyl)-1,1-(ethylenedioxy)-7-[1'-(ethoxy-carbonyl)propyl]-5-oxo-Δ⁶⁽⁸⁾-tetrahydroindolizine
(Compound 7).

A solution of the ester ketal 6 (2.0 g, 6.0 mmol) in acetic anhydride (30 mL) and HOAc (10 mL) was hydrogenated for 6 hr at 45°C under 50 psi in the 15 presence of Raney nickel (3 g; washed with HOAc). The catalyst was removed by filtration and the solvent removed in vacuo to give 7 (2.3 g, 100%) as an oil. Purification by silica gel column chromatography (2% MeOH in CHCl₃) gave pure 7 as an oil.

20 6-(Acetoxymethyl)-1,1-(ethylenedioxy)-7-[1'-(ethoxy-carbonyl)propyl]-5-oxo-Δ⁶⁽⁸⁾-tetrahydroindolizine
(Compound 9).

A cooled solution of amide 7 (2.3 g, 6.0 mmol) in Ac₂O (30 mL) and HOAc (10 mL) was treated with NaNO₂ 25 (1.8 g, 26 mmol) and the reaction mixture stirred for 2 h at 0°C. Inorganic salts were removed by filtration and the solvent removed in vacuo at room temperature to afford the N-nitroso intermediate 8 as an oil. Compound 8 was converted directly to the title acetoxo compound 9 30 by refluxing overnight in CCl₄. The solution was washed with water, dried (Na₂SO₄) and the solvent removed in vacuo to give 9 (2.3 g 100%) as an oil.

1,1'Ethylenedioxy-5-oxo-(5'-ethyl-5'-hydroxy-2'H,5'H,6'H-6-oxopyrano)-[3',4'-f]-Δ^{6,8}-tetrahydro-

indolizine (Compound 10).

Oxygen was bubbled through a mixture of 6-(acetoxymethyl)-1,1-(ethylenedioxy)-7-[1'-ethoxy-carbonyl]-propyl]-5-oxo- $\Delta^{6,8}$ -tetrahydroindolizine 5 (Compound 9, 405 mg, 1.07 mmol), anhydrous K_2CO_3 (148 mg 1.07 mmol) and methanol (7.5 mL) for 24 hr. The solution was cooled in an ice bath and made acidic (pH 2-4) by addition of 1N H_2SO_4 . Most of the methanol was removed in vacuo at room temperature, and water (20 mL) was 10 added. The aqueous solution was extracted with CH_2Cl_2 (3 x 20 mL), dried ($NaSO_4$) and evaporated to give a solid which was crystallized from CH_2Cl_2 -hexane to give 280 mg (85%) of 10: mp 179-181°C; ν_{max} ($CHCl_3$) 1740, 1660 cm^{-1} ; 1H -NMR ($CDCl_3$) δ 0.91 (t, 3, J=7 Hz, CH_2CH_3), 1.75 (q, 2, 15 J=7 Hz, CH_2CH_3), 2.35 (t, 2, J=6.5 Hz, CH_2 α to ketal), 4.1 (m, 6, OCH_2CH_2O and CH_2N), 5.30 (m, 2, $ArCH_2O$), 6.87 (s, 1, pyridone). Anal. Calcd for $C_{15}H_{17}NO_6$: C, 58.63; H, 5.54; N, 4.56. Found: C, 58.72, H, 5.68; N, 4.57.

5'RS-1,5-Dioxo-(5'-ethyl-5'-hydroxy-2'H,5'H,6'H- 20 6-oxopyrano)-[3',4',f]- $\Delta^{6,8}$ -tetrahydroindolizine (Compound 11).

A solution of 10 (3.88 g, 12.6 mmol) in 2N H_2SO_4 (50 mL) and DME (50 mL) was heated for 24 hr under N_2 . The reaction mixture was concentrated to one half its 25 volume in vacuo, diluted with H_2O (100 mL) and extracted with CH_2Cl_2 (5 x 50 mL). The organic layer was dried (Na_2SO_4) and evaporated to yield a solid which was crystallized from CH_2Cl_2 -hexane to yield 2.68 g (80%) of 11 as a light brown solid: mp 185-187°C; ν_{max} ($CHCl_3$) 30 1750 (shoulder, ketone), 1745 (lactone), 1660 cm^{-1} (pyridone); 1H -NMR ($CDCl_3$) δ 0.91 (t, 3, J=7 Hz, CH_2CH_3), 1.80 (q, 2, J=7 Hz, CH_2CH_3), 2.93 (t, 2, J=6.5 Hz, $CH_2C=O$), 4.30 (t, 2, J=6.5 Hz, CH_2N), 5.35 (m, 2, $ArCH_2O$), 7.17 (s, 1, aromatic H). Anal. Calcd for 35 $C_{13}H_{13}NO_5$: C, 59.32; H, 4.94; N, 5.32. Found: C, 59.12, H, 4.91; N, 5.16.

SYNTHESIS OF CAMPTOTHECIN ANALOGS:Synthesis of 11-hydroxy-20(RS)-camptothecin.

11-hydroxy-20(RS)-camptothecin is prepared from 11-methoxy-20(RS)-camptothecin by demethylation of the latter with hydrobromic acid as follows:

11-Methoxy-20(RS)-camptothecin.

A mixture of 4-methoxy-2-aminobenzaldehyde (180 mg, 1.19 mmol) and the tricyclic ketone 11 (300 mg, 1.14 mmol) in toluene (18 mL) was heated under N₂ in a flask equipped with a Dean-Stark trap. At reflux p-toluenesulfonic acid (5 mg) was added, and the red-brown solution was heated for an additional 2 hr. The toluene was removed under reduced pressure to give a brown solid which was treated with water (10 mL) and chloroform (20 mL). The aqueous phase was extracted with additional chloroform (3 × 20 mL) and the combined extracts dried (Na₂SO₄). Evaporation gave a brown solid which was recrystallized from methanol-chloroform to give 216 mg (50%) of compound as a tan solid: 275-279°C; mass spectrum (electron impact), m/z 378.1219 M⁺; C₂₁H₁₈N₂O₅ requires 378.1214; ν_{max} (KBr) 3480 (OH), 1745 (lactone), 1660 (pyridone), 1622, 1236 and 1152 cm⁻¹; ¹H-NMR (DMSO-d₆) δ 0.87 (t, 3, J=7 Hz, H-18), 1.85 (m, 2, H-19), 3.95 (s, 3, 11-OCH₃), 5.24 (s, 2, H-5), 5.42 (s, 2, H-17), 7.32 (s, 1, H-14), 7.37 (dd, 1, J = 9, 2.5 Hz, H-10), 7.56 (d, 1, J = 2.5 Hz, H-12), 8.02 (d, 1, J=9 Hz, H-9), 8.60 (s, 1, H-7).

11-Hydroxy-20(RS)-camptothecin.

11-methoxy-20(RS)-camptothecin (75 mg) was combined with 48% aqueous HBr (2.5 mL) and heated at reflux for 6 hr. The red-brown mixture was stripped of solvent under high vacuum. Chromatography of the residue through silica gel (15 g) (7% MeOH-CHCl₃) gave the 11-hydroxy compound (33 mg, 45%) which was further purified by

recrystallization from 13% MeOH in CHCl_3 : mp 323-326°C;
mass spectrum (electron impact), m/z 364.1054 M^+ ,
 $\text{C}_{20}\text{H}_{16}\text{N}_2\text{O}_5$ requires 364.1059; ν_{max} (KBr) 3450, 1742,
1654, 1613, 1592, 1570, 1245 cm^{-1} ; λ_{max} (EtOH), 224
5 (log ϵ 4.58), 259, (4.39), 353 (4.16), 371 (4.19), 387
(4.20); $^1\text{H-NMR}$ (DMSO-d_6): δ 0.88 (t, 3, $J=7$ Hz, H-18),
1.85 (m, 2, H-19), 5.20 (s, 2, H-5), 5.41 (s, 2, H-17),
6.51 (br s, 1, OH-20), 7.26 (dd, 1, $J=9$, 2.5 Hz, H-10),
7.28 (s, 1, H-14).

10 10-Hydroxy-20(RS)-camptothecin.

This compound is prepared in a manner analogous to
that described for the 11-hydroxycamptothecin using
5-methoxy-2-aminobenzaldehyde which is reacted with the
tricyclic ketone 11 in the presence of p-toluenesulfonic
15 acid. The product is 10-methoxy-20(RS)-camptothecin
which on treatment with refluxing hydrobromic acid as
described for 11-hydroxycamptothecin, gives 10-hydroxy-
20(RS)-camptothecin.

20 9-Methoxy-20(RS)-camptothecin and 9-Hydroxy-
20(RS)-camptothecin

In a manner analogous to that described for
11-methoxy-20(RS)-camptothecin, 6-methoxy-2-aminobenz-
aldehyde is treated with the tricyclic 11 ketone in the
presence of p-toluenesulfonic acid yielding 9-methoxy-
25 20(RS)-camptothecin. Demethylation with hydrobromic acid
gives 9-hydroxy-20(RS)-camptothecin.

10-Nitro-20(RS)-camptothecin.

A mixture of 2-amino-5-nitrobenzaldehyde (95 mg,
0.57 mmol) and the tricyclic ketone 11 (150 mg,
30 0.57 mmol) was heated at 120°C for 10 min. The
temperature was raised to 160°C, and the dark molten mass
was kept at this temperature for 1.5 hr with occasional
stirring. Chromatography of the residue through silica
gel (20 g) using 0.5% MeOH in CHCl_3 afforded the title

compound (108 mg) as a yellow solid; mp 297-300°C (decomp.); mass spectrum (electron impact), m/z 393.0965 M^+ , $C_{20}H_{15}N_3O_6$ requires 393.0960; ν_{\max} (KBr) 3450 (OH), 1745 (lactone), 1660 (pyridone), 1620, 1350, 5 and 1160 cm^{-1} ; 1H -NMR (TFA- d_1) δ 1.14 (t, 3, $J=7$ Hz, H-18), 2.15 (m, 2, H-19), 5.88 (s, 2, H-5), 5.68 (Abq, 2, $J=17$ Hz, $\Delta\gamma = 85$ Hz, H-17), 8.43 (s, 1, H-14), 8.70 (d, 2, $J=8$ Hz, H-12), 9.05 (d, 2, $J=8$ Hz, H-11), 9.35 (s, 1, H-9), 9.60 (s, 1, H-7).

10 10-Amino-20(RS)-camptothecin

A suspension of 10-nitro-20(RS)-camptothecin (100 mg) and 10% Pd/C (40 mg) in absolute EtOH (40 mL) was stirred in an atmosphere of H_2 at room temperature for 30 min. Filtration through Celite and removal of the solvent under reduced pressure gave a tan yellow solid (86 mg crude). Recrystallization from 13% MeOH/ $CHCl_3$ gave the pure product (30 mg) as an olive-yellow solid; mp, softening at 135°C, gradual blackening upon further heating; mass spectrum (electron impact), m/z 363.1116 M^+ ; 20 $C_{20}H_{17}N_3O_4$ requires 363.1218; ν_{\max} (KBr) 3440 (OH, NH_2), 1750 (lactone), 1660 (pyridone) cm^{-1} ; 1H -NMR (TFA- d_1) δ 1.06 (t, 3, $J = 7$ Hz, H-18), 2.08 (d, $J = 7$ Hz, H-17), 5.89 (s, 2, H-5), 5.70 (Abq, 2, $J = 17$ Hz, $\Delta\gamma = 85$ Hz, H-17), 8.34 (d, $J = 9$ Hz, H-12), 8.64 (d, $J = 9$ Hz, H-11), 25 9.26 (s, 1, H-9), 9.43 (s, 1, H-7).

9-Nitro-20(RS)-camptothecin and 9-Amino-20(RS)-camptothecin.

A mixture of 2-amino-6-nitrobenzaldehyde is treated with the tricyclic ketone 11 in the manner described for 30 the 10-nitro series above yielding 9-nitro-20(RS)-camptothecin. This compound, after reduction with palladium/carbon, yielded 9-amino-20(RS)-camptothecin. Alternatively, the 9-amino compound is obtained in one step by reaction of 2,6-diaminobenzaldehyde with ketone 35 11.

11-Nitro-20(RS)-camptothecin and 11-Amino-20(RS)-camptothecin.

In a manner similar to that described for 10-nitro-20(RS)-camptothecin, a mixture of 2-amino-4-nitrobenzaldehyde is treated with the tricyclic ketone 11 yielding 11-nitro-20(RS)-camptothecin which in turn is reduced to 11-amino-20(RS)-camptothecin by palladium/carbon. Alternatively, the 11-amino-20(RS)-camptothecin is obtained by reaction of 2,4-diaminobenzaldehyde with ketone 11.

10,11-Dihydroxy-20(RS)-camptothecin.

A solution of the crude dibenzylxy aminoacetal (400 mg) and the tricyclic ketone 11 (132 mg, 0.5 mmol) in toluene (60 mL) was refluxed for 8 hr. It was filtered hot, and the pure dibenzylether was collected upon cooling (200 mg, 81%); mp 276°C. ν_{\max} (KBr) 3440, 1740, 1650, 1590, 1490, 1440, 1380, 1250, 1140, 1100 cm^{-1} ; 250 MHz $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ 0.88 (t, 3, $J = 7 \text{ Hz}$, H-18), 1.86 (m, 2, H-19), 5.22 (s, 2, H-17), 5.34 (s, 2, 10- $\text{OCH}_2-\text{C}_6\text{H}_5$), 5.39 (s, 2, 11- $\text{OCH}_2-\text{C}_6\text{H}_5$), 5.41 (s, 2, H-5), 6.5 (s, 1, OH), 7.25 (s, 1, H-14) 7.35-7.65 (m, 12, H-9, 12, $-\text{OCH}_2-\text{C}_6\text{H}_5$), 8.44 (s, 1, H-7). Anal. calcd for $\text{C}_{34}\text{H}_{28}\text{N}_2\text{O}_6$: C, 72.84; H, 5.03; N, 5.00. Found C, 72.91; H, 5.09; N, 4.96.

The dibenzyl ether (130 mg, 0.23 mmol) was mildly refluxed for 2 hr in 24% HBr (50 mL). The acid was removed, and the residue was dissolved in hot methanol (50 mL). Ether (50 mL) was added at room temperature and the yellow powdery dihydroxy camptothecin hydrobromide was collected (122 mg, 77%) mp > 300°C. ν_{\max} (KBr) 3400 (b), 1740, 1655, 1585, 1545, 1510, 1395, 1300, 1270, 1200, 1160 cm^{-1} ; $^1\text{H NMR}$ (DMSO, d_6): δ 0.88 (t, 3, $J=7 \text{ Hz}$, H-18), 1.85 (m, 2, H-19), 5.20 (s, 2, H-17), 5.42 (s, 2, H-5), 7.31 (s, 2, H-9, H-14), 7.40 (s, 1, H-12), 8.45 (s, 1, H-7). Anal. calcd for $\text{C}_{20}\text{H}_{17}\text{BrN}_2\text{O}_6$

· 0.5 H₂O: C, 51.08; H, 3.86; N, 5.95; Br, 16.99. Found C, 51.09; H, 4.04; N, 5.78; Br, 16.83.

Dihydroxy hydrobromide salt (110 mg, 0.23 mmol) was suspended in water (10 mL). Sodium hydroxide (0.1 N, 7.2 mL) was added and the mixture was agitated. The resulting clear solution was acidified using 5N HCl; and after an hour, the sample was centrifuged, the supernatant liquid was decanted and the process repeated with additional water (20 mL). The residue was dried (78 mg, 74%); mp > 300°C. ν_{\max} (KBr): 3490, 3000 (b), 1740, 1645, 1590, 1460, 1385, 1265, 1190, 1150 cm⁻¹. ¹H NMR (DMSO, d₆): δ 0.88 (t, 3, J=7 Hz, H-18), 1.87 (q, 2, H-19), 5.20 (s, 2, H-17), 5.42 (s, 2, H-5), 7.35 (s, 1, H-14), 7.44 (s, 1, H-9), 7.52 (s, 1, H-12), 8.51 (s, 1, H-7). Anal. calcd for C₂₀H₁₆N₂O₆ · 0.75 H₂O: C, 61.06; H, 4.44; N, 7.12. Found C, 61.12; H, 4.44; N, 7.09.

10-Chloro-20(RS)-camptothecin.

This compound was prepared by treating 5-chloro-2-aminobenzaldehyde with the tricyclic ketone 11.

A solution of the 5-chloro-2-aminobenzaldehyde (80 mg, 0.51 mmol) and the tricyclic ketone 11 (100 mg, 0.38 mmol) in toluene (60 mL) was refluxed for 15 min. p-Toluenesulfonic acid (10 mg) was then added, and refluxing was continued for an additional 5 hr. The solvent was removed in vacuo and the residue chromatographed (silica gel 60, 2% MeOH-CHCl₃). The product obtained was recrystallized from CHCl₃-MeOH-EtOAc; mp 270°C, 60 mg (41%). ν_{\max} (KBr), 3430, 1745, 1655, 1600, 1495, 1230, 1160 cm⁻¹. 250 MHz ¹H-NMR (TFA-d₁) δ 1.15 (t, 3, J=7 Hz, H-18), 2.16 (m, 2, H-19), 5.73 (ABq, 2, J=17 Hz, $\Delta\gamma$ = 85 Hz, H-17), 5.84 (s, 2, H-5), 8.29 (d, 1, J=9 Hz, H-11), 8.35 (s, 1, H-14), 8.40 (s, 1, H-9), 8.45 (d, 1, J=9 Hz, H-12), 9.31 (s, 1, H-7). Anal. calcd for C₂₀H₁₅ClN₂O₄ 0.5 H₂O: C, 61.47; H, 4.12; N, 7.17; Cl, 9.07. Found C, 61.41; H, 4.12; N, 7.12;

Cl, 9.11.

10-Methyl-20(RS)-camptothecin.

5-Methyl-2-aminobenzaldehyde was treated with the tricyclic ketone 11 to give the title compound.

- 5 The tricyclic ketone 11 (130 mg, 0.5 mmol) and the 5-methyl-2-aminobenzaldehyde (560 mg) in toluene (60 mL) were refluxed for 0.5 hr. Acetic acid (1 mL) and p-toluenesulfonic acid (35 mg) were added, and refluxing was continued for an additional 5 hr. The solvent was removed in vacuo, and the residue was triturated with warm ether (30 mL). The product was recrystallized from chloroform-methanol-ether to yield pure compound (102 mg, 57%), mp 278-280°C. (KBr) 3460, 2980, 1740, 1655, 1590, 1550, 1470, 1450, 1370, 1260, 1240, 1160, 1050 cm⁻¹.
15 250 MHz ¹H-NMR (DMSO-d₆) δ 0.89 (t, 3, J=7 Hz, H-18), 1.87 (q, 2, H-19), 2.54 (s, 3, 10-CH₃), 5.24 (s, 2, H-17), 5.42 (s, 1, H-5), 7.31 (s, 1, H-14), 7.69 (d, 1, J=8.6 Hz, H-11), 7.86 (s, 1, H-9), 8.05 (d, 1, J=8.6 Hz; H-12), 8.55 (s, 1, H-7). Anal. calcd for
20 C₂₁H₁₈N₂O₄ · 0.25 H₂O: C, 68.75; H, 5.08; N, 7.64. Found C, 68.74; H, 5.08; N, 7.64.

11-Formyl-20(RS)-camptothecin.

- 2-Nitroterephthaldicarboxaldehyde was converted to the ethylene diacetal by conventional methods and reduced
25 using Na₂S. A solution of the nitro diacetal (4.1 g, 17.5 mmol), Na₂S (14 g) in 80% ethanol (15 mL) was refluxed for 1 hr. Ethanol was removed in vacuo, the reaction mixture was diluted with water (10 mL) and the aqueous phase was extracted with CH₂Cl₂ (4 · 50 mL). The
30 organic phase was washed with water, dried (MgSO₄), and evaporated to give the aminodiacetal, which was recrystallized from ethyl acetate-hexane (2.8 g, 78%); mp 76°C. ν_{max} (KBr) 3480, 3395, 3000, 2960, 2900, 1625, 1445, 1395, 1085, 950 cm⁻¹. 60 MHz ¹H NMR
35 (CDCl₃-D₂O) δ 4.0 (m, 8, -OCH₂CH₂O-), 5.6 (s, 1,

-O-CH₂-O-, C-4), 5.7 (s, 1, -O-CH₂-O-, C-1), 6.6 (s, 1, H-3), 6.65 (d, 1, J=8 Hz, C-5), 7.2 (d, 1, J=8 Hz, H-6). Anal. calcd for C₁₂H₁₅NO₄: C, 60.66; H, 6.36; N, 5.90. Found C, 60.79; H, 6.41; N, 5.84.

- 5 A solution of the tricyclic ketone 11 (265 mg, 1.0 mmol), aminodiacetal (500 mg, 2.1 mmol), 300 mg initially, 100 mg each at intervals of 5 and 10 hr) in toluene (70 mL) was refluxed for 0.5 hr. Acetic acid (2 mL) was added and refluxing continued for 18 hr. The
10 solvent was evaporated in vacuo, and the residue was taken up in 75% methanol (250 mL). Conc. HCl (3 mL) was added and the reaction mixture heated at 50-60°C for 24 hr. The mixture was filtered, and the residue was washed with water and recrystallized from
15 CHCl₃-MeOH-EtOAc. mp: 276-279°C (175 mg, 45%). ν_{\max} (KBr) 3460, 1745, 1690, 1655, 1600, 1200, 1150, 1135 cm⁻¹. 250 MHz ¹H NMR (TFA-d₁), δ 1.16 (t, 3, J=7 Hz, H-18) 2.16 (q, 2, J=7 Hz, H-19), 5.78 (ABq, 2, J=18 Hz, $\Delta\gamma$ = 85 Hz, H-17), 5.89 (s, 2, H-5), 8.43 (s, 1, H-14), 8.66 (d, 1, J=8.5 Hz, H-10), 8.60 (d, 1, J=8.5 Hz, H-9), 9.12 (s, 1, H-12), 9.49 (s, 1, H-7), 10.42 (s, 1, CHO). Anal. calcd. for C₂₁H₁₆N₂O₅ · H₂O: C, 64.01; H, 4.56; N, 7.11. Found C, 64.08, H, 4.21; N, 6.84.

11-Cyano-20(RS)-camptothecin.

- 25 A mixture of 11-formyl-20(RS)-camptothecin (225 mg, 0.6 mmol), hydroxylamine hydrochloride (50 mg, 0.72 mmol), sodium formate (90 mg, 1.3 mmol), and formic acid (6 mL) was refluxed for 1.5 hr. The mixture was evaporated to dryness in vacuo, and the residue was
30 washed with water, dried and chromatographed (silica gel 60, 0.5% MeOH-CHCl₃) and recrystallized from CHCl₃-EtOAc to yield the 11-cyano compound (65 mg, 29%): mp 288°C. ν_{\max} (KBr) 3400, 2235, 1735, 1655, 1590, 1450, 1400, 1230, 1150, 1110, 1045 cm⁻¹. 250 MHz ¹H NMR (DMSO-d₆):
35 δ 0.88 (t, 3, J=7 Hz, H-18), 1.88 (m, 2, H-19), 5.32 (s, 2, H-17), 5.44 (s, 2, H-5), 7.37 (s, 1, H-14), 7.98

(d, 1, J=8.5 Hz, H-10), 8.32 (d, 1, J=8.5 Hz, H-9), 8.74 (s, 1, H-12), 8.80 (s, 1, H-7). Anal. calcd for $C_{21}H_{15}N_3O_4 \cdot 1.5 H_2O$: C, 62.99; H, 4.52; N, 10.49. Found C, 62.99; H, 3.95; N, 10.20.

- 5 Alternatively, 11-cyano-20(RS)-camptothecin can be prepared by the reaction of 5-cyano-2-aminobenzaldehyde with the tricyclic ketone 11.

PREPARATION OF CAMPTOTHECIN ANALOGS WITH MODIFIED A RING STRUCTURE:

- 10 The reaction of the tricyclic ketone 11 with suitable precursors other than substituted ortho-amino-benzaldehydes can be used to give active new camptothecin analogs exemplified by the following non-limiting examples:

15 10-Aza-20(RS)-camptothecin.

- A solution of 4-aminonicotinaldehyde (24.2 mg, 0.198 mmol), the tricyclic ketone 11 (53.5 mg, 0.203 mmol) and p-TsOH \cdot H₂O (2 mg) in toluene (25 mL) was refluxed for 4 days using Dean-Stark trap. The
20 solvent was removed under reduced pressure, and the residue was chromatographed through silica gel (20 g) using CHCl₃-acetone-MeOH (5:1:1). The product was crystallized from 13% MeOH in CHCl₃ and EtOAc: mp 289-292°C; mass spectrum (electron impact), m/z 349.1061 M⁺;
25 C₁₉H₁₅N₃O₄ requires 349.1066; ν_{\max} (KBr) 3320 (OH), 1730 (lactone), 1650 (pyridone), 1600 (aromatic) cm⁻¹; ¹H NMR (CDCl₃) 1.05 (t, 3, J=7.3 Hz, H-18), 1.92 (m, 2, H-19), 5.35 (s, 2, H-5), 5.52 (ABq, 2, J=18 Hz, $\Delta\gamma$ = 85 Hz, H-17), 7.74 (s, 1, H-14), 8.04 (d, 1, J=5.5 Hz, H-12),
30 8.53 (s, 1, H-7), 8.84 (d, J=5.5 Hz, H-11), 9.4 (s, 1, H-9).

A-Nor-9-thia-20(RS)-camptothecin.

This sulfur containing camptothecin analog is prepared by the reaction of 3-amino-2-formylthiophene

with tricyclic ketone 11.

A solution of 3-amino-2-formylthiophene (79 mg, 0.62 mmol) and the tricyclic ketone 11 (96 mg, 0.37 mmol) in toluene (1.5 mL) was brought to reflux and then cooled before adding a crystal of p-toluenesulfonic acid. The mixture was refluxed for 2.5 hr under N₂, cooled and the precipitate filtered. The crude material was chromatographed on silica gel (20 g) by elution with 2% MeOH in CHCl₃. Crystallization of the product from 13% MeOH-CHCl₃ and EtOAc yielded the title compound as a yellow solid (19 mg, 15%): mp 297-298°C; ν_{\max} 1740 (lactone), 1655 cm⁻¹ (pyridone); ¹H NMR (TFA-d₁) δ 1.05 (t, 3, J=7 Hz, H-18), 2.07 (q, 2, J=7 Hz, H-19), 5.60 (m, 2, H-17), 5.65 (s, 2, H-5), 7.89 (d, J=6 Hz, H-11), 8.05 (s, 1, H-14), 8.57 (d, J=6 Hz, H-10), 9.23 (s, 1, H-7). Anal. (C₁₈H₁₄N₂O₄S), calcd. C, 61.02; H, 3.95; N, 7.91. Found C, 60.65; H, 4.01; N, 7.78.




10,11-Methylenedioxy-20(RS)-camptothecin.

The required ortho-aminoaldehyde was prepared by reduction of 2-nitropiperonal. This compound (60 mg, 0.36 mmol) and the tricyclic ketone 11 (53 mg, 0.20 mmol) were refluxed for 8 hr in toluene (30 mL) containing p-TsOH · H₂O (8 mg). The solvent was removed *in vacuo*, the red residue adsorbed onto Celite (1 g) and chromatographed through silica gel (10 g) using 3% MeOH in CHCl₃. Concentration of the appropriate fractions gave 10,11-methylenedioxy-20(RS)-camptothecin (36 mg, 45%) as a pale tan solid. Crystallization of this material from CHCl₃ gave the analytical sample as a cream-colored solid: mp > 250°C (decomp); ν_{\max} (KBr) 1750 (lactone), 1655 (pyridone), 1585 cm⁻¹ (aromatic); ¹H NMR (TFA-d₁) δ 1.15 (t, 3, J=7 Hz, H-18), 2.16 (q, 2, J=7 Hz, H-19), 5.76 (ABq, 2, J=17 Hz, $\Delta\gamma$ = 85 Hz, H-17), 5.73 (s, 2, H-5), 6.44 (s, 2, OCH₂O), 7.55 (s, 1, H-14), 7.69 (s, 1, H-9), 8.16 (s, 1, H-12), 9.05 (s, 1, H-7). Anal. calcd for C₂₁H₁₆N₂O₆:


392.1008. Found 392.1009 ($C_{21}H_{16}N_2O_6 \cdot 1.0 H_2O$).

PREPARATION OF WATER SOLUBLE CAMPTOTHECIN ANALOGS:


The following non-limiting examples of water soluble camptothecin analogs were prepared as follows. The 9-
5 glycinamido and 9-succinamido, 9-(N-methylpiperazinocarbonylamino), 9-(N,N-diethylaminoethoxy carbonylamino), and camptothecin analogs have the structures shown below.




3. N-Methylpiperazinocarbonylamino,



4. N,N-Diethylaminoethoxycarbonylamino,



5. Diethylaminoethoxy,



Synthesis of 9-Glycinamido-20(RS)-camptothecin
Hydrochloride

9-Amino-20(RS)-camptothecin was prepared from 2-amino-6-nitrobenzaldehyde and tricyclic ketone 11 by the
5 process described above.

9-(tert-Butoxycarbonylglycinamido)-2-(RS)-camptothecin.

A stirred solution of 9-amino-20(RS)-camptothecin (88 mg, 0.242 mmol) and N-(tert-butoxycarbonyl)glycine (110 mg, 0.629 mmol) in dry N,N-dimethylformamide (10 ml)
10 under nitrogen was treated with dicyclohexylcarbodiimide (125 mg, 0.607 mmol) at room temperature. After stirring for 18 hr., the turbid white mixture was filtered to remove the dicyclohexylurea byproduct. The solvent was removed by high vacuum distillation, and the tan-yellow
15 residue was chromatographed through silica gel (25 g) using a stepwise gradient of 250 ml each of chloroform, 1% methanol/chloroform, and 2% methanol/chloroform. The appropriate fractions afforded 55 mg (44%) of the 9-tert-butoxycarboxyl glycinamido compound as a yellow solid.
20 Recrystallization from methanol provided the sample as a beige solid, mp 208-210°C. ν_{\max} (KBr) 3360 (br, OH, amide NH), 1750 (lactone), 1710 (carbamate), 1692 (amide), 1660 (lactone), 1622, 1598, 1493, 1370, 1256, 1235, 1165, 1110, 1058, 1032, 825 and 728 cm^{-1} ; ^1H NMR
25 (DMSO- d_6) δ 0.89 (t, 3, J=7Hz, H-18), 1.44 (s, 9, C(CH₃)₃), 1.88 (m, 2, H-19), 3.92 (d, 2, J=6Hz, COCH₂N), 5.29 (s, 2, H-5), 5.44 (s, 2, H-17), 6.53 (s, 1, OH), 7.19 (t, 1, J=6Hz, CH₂NHCO), 7.37 (s, 1, H-14), 7.79 (d, 1, J=7Hz, H-10), 7.85 (t, 1, J=7Hz, H-11), 8.03 (d, 1, J=7Hz, H-12), 8.79 (s, 1, H-7), and 10.20 (s, 1, amide
30 H). Anal. calcd. for C₂₇H₂₈N₄O₇·H₂O: C, 60.21; H, 5.61; N, 10.40. Found: C, 60.35; H, 5.64; N, 10.23.

9-Glycinamido-20(RS)-camptothecin Hydrochloride.

The tert-butoxycarboxylglycinamide derivative from
35 above (21 mg, 0.040 mmol) was suspended in methylene chloride (10 ml) under nitrogen and then dissolved by the

addition of methanol (0.75 ml). The stirred solution was chilled to 0°C and treated over 5 min. with a saturated solution of hydrogen chloride in anhydrous dioxane (4.5 ml) resulting in a turbid yellow solution. The stirred mixture was left to warm to room temperature, and after 2 hr. the solvents were removed under reduced pressure to give the deprotected compound as an orange-yellow solid (18 mg). The sample was taken up in deionized water (5 ml) and the hazy yellow solution filtered through a 0.45 μ m membrane filter to remove extraneous water-insoluble material. The clear filtrate was lyophilized to provide the pure salt as a fluffy yellow solid (14 mg, 77%), mp darkening above 245°C with no melting up to 310°C. ν_{max} (KBr) 2400-3650 cm^{-1} (OH, amide H, amine HCl salt), 1742 (lactone), 1700 (amide), 1658 (pyridone), 1590, 1550, 1495, 1234, 1163, 1110, 1050, 902, 820 and 720 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 0.89 (t, J=7.5Hz, H-18), 1.89 (m, 2, H-19), 4.03 (d, 2, J=5.4Hz, COCH_2N), 5.30 (s, 2, H-5), 5.44 (s, 2, H-17), 7.37 (s, 1, H-14), 7.86 (d, 1, J=7Hz, H-12), 7.92 (t, 1, J=7Hz, H-11), 8.07 (d, 1, J=7Hz, H-10), 8.35 (br s, 3, NH_3^+), 8.95 (s, 1, H-7), 10.88 (s, 1, amide H). Anal. calcd. for $\text{C}_{22}\text{H}_{21}\text{ClN}_4\text{O}_5 \cdot 3\text{H}_2\text{O}$: C, 51.71; H, 5.32; Cl, 6.94; N, 10.96. Found: C, 51.82; H, 5.23; Cl, 6.75; N, 10.61.

Synthesis of 9-Succinamido-20(RS)-camptothecin, Sodium Salt.

The 9-succinamido derivative is synthesized from 9-amino-20(RS)-camptothecin (synthesis described above) by the following method.

9-Succinamido-20-(RS)-camptothecin.

A stirred suspension of 9-amino-20(RS)-camptothecin (400 mg, 1.102 mmol) and succinic anhydride (125 mg, 1.25 mmol) in pyridine (5 ml) under nitrogen was heated at 95°C for 2 hr. The solvent was removed from the brown solution by high vacuum distillation to give the crude

amide as a brown gum. Purification was effected by chromatography through silica gel employing a solvent gradient from 1 : 5% methanol/chloroform to 1 : 50% methanol/chloroform. Evaporation of the appropriate fractions gave 272 mg of the 9-succinamido compound as an orange-tan solid (53%), and recrystallization from methanol gave the material as a light tan solid, mp 265-270°C (decomp). ν_{\max} (KBr) 2600-3650 (OH, NH, acid), 1738 (lactone), 1650 (pyridone), 1520-1610 (broad), 1485, 1400, 1232, 1160, 1110, 1050, 818, 720 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 0.89 (t, 3, $J=7\text{Hz}$, H-18), 1.88 (m, 2, H-19), 2.45-2.50 (m, 4, $-\text{NCOCH}_2\text{CH}_2\text{CO}_2\text{H}$), 5.24 (s, 2, H-5), 5.42 (s, 2, H-17), 6.53 (br s, 1, OH), 7.33 (s, 1, H-14), 7.77 (t, 1, $J=7\text{Hz}$, H-11), 7.85 (d, 1, $J=7\text{Hz}$, H-12), 7.91 (d, 1, $J=7\text{Hz}$, H-10), 8.83 (s, 1, H-7), 10.73 (br s, 1, amide H or CO_2H). Calcd. for ($\text{M}^+-\text{CO}_2-\text{H}_2\text{O}$): 401.1375. Found: 401:1368. Calcd. for $\text{C}_{24}\text{H}_{21}\text{N}_3\text{O}_7 \cdot 2.5\text{H}_2\text{O}$: C, 56.68; H, 4.73; N, 8.45. Found: C, 56.69; H, 4.65; N, 8.26.

Alternatively, the 9-succinamido derivative can be prepared by hydrolysis of its ethyl ester which is prepared by the following general method:

9-Amino-20(RS)-camptothecin in dry N,N-dimethylformamide containing pyridine is reacted at 0-10°C with a slight excess of ethylsuccinyl chloride in N,N-dimethylformamide solution. After work-up and chromatography on silica gel, a 75% yield of the 9-(ethyl)glycinamide derivative is obtained.

9-Succinamido-20(RS)-camptothecin, Sodium Salt.

The preceding succinamide free acid (151 mg, 0.326 mmol) was suspended in methanol (5 ml) and the stirred mixture treated dropwise at room temperature over 5 min with 0.1 N aqueous sodium hydroxide solution (3.26 ml, 0.326 mmol, 1 eq.). During the addition the solution became hazy yellow-orange, and at the end of the addition the pH was near 7. The methanol was evaporated under reduced pressure, and the resulting aqueous solution was

diluted with deionized water (6 ml). Filtration through a 0.5 μ m membran filter was followed by lyophilization to provide the 9-succinamido-20(RS)-camptothecin salt as a fluffy orange-yellow solid (145 mg, 92%), mp >200°C (decomp). ^1H NMR (DMSO- d_6): δ 0.89 (t, 3, J=7Hz, H-18), 1.88 (m, 2, H-19), 2.41-2.60 (m, 4, -NCOCH₂CH₂CO₂Na), 5.30 (s, 2, H-5), 5.43 (s, 2, H-17), 6.55 (br s, 1, OH), 7.33 (s, 1, H-14), 7.76 (t, 1, J=8Hz, H-11), 7.87 (d, 1, J=8Hz, H-12), 8.16 (d, 1, J=8Hz, H-10), 12.51 (br s, 1, amide H).

Synthesis of 9-Diethylaminoethoxy-20(RS)-camptothecin Hydrochloride

The title compound is prepared from 9-hydroxy-20(RS)-camptothecin (synthesis of hydroxy-camptothecins described on pp. 30-32 of current Patent Application) in the following manner:

9-Diethylaminoethoxy-20(RS)-camptothecin.

A stirred mixture of 9-hydroxy-20(RS)-camptothecin (20 mg, 0.055 mmol), N,N-diethylaminoethylchloride hydrochloride (15.4 mg, 0.090 mmol), and powdered anhydrous potassium carbonate (30 mg, 0.276 mmol) in anhydrous dimethylformamide (0.5 ml) was heated under nitrogen at 55°C for 3 h. During the reaction, the mixture became clear yellow-orange and then tan. The solvent was removed by high vacuum distillation, and the tan residue was dispersed on Celite and chromatographed through silica gel (5 g) using 10% methanol/chloroform. Evaporation of the appropriate fractions gave 14 mg (55%) of the desired aminoether as a pale yellow solid. Recrystallization from ethyl acetate afforded the pure title compound as a beige solid, mp 173-176°C (decomp). γ_{max} (KBr) 3150-3650 (br. OH), 2967 and 2920 (CH), 1745 (lactone), 1658 (pyridione), 1592-1620 (aromatic), 1469, 1460, 1384, 1370, 1267, 1232, 1190, 1157, 1110, 1050, 810 and 720 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 0.89 (t, 3, J=7 Hz, H-18), 1.11 (t, 6, J=7 Hz, -N(CH₂CH₃)₂), 1.89 (m, 2, H-19),

2.82 (br s, 4, $-\text{N}(\text{CH}_2\text{CH}_3)_2$; with D_2O exchange signal is at 2.88 δ as a quartet, $J=7$ Hz), 3.17 (br s, 2, $-\text{OCH}_2\text{CH}_2\text{NET}_2$; with D_2O exchange, signal is at 3.24 δ as a fine triplet), 4.37 (m, 2, $-\text{OCH}_2\text{CH}_2\text{NET}_2$), 5.29 (s, 2, H-5), 5.43 (s, 2, H-17), 6.53 (s, 1, 20-OH), 7.19 (d, 1, $J=7$ Hz, H-10), 7.33 (s, 1, H-14), 7.76 (m, 2, H-11 and H-12), 8.87 (s, 1, H-7). Anal. calcd. for $\text{C}_{26}\text{H}_{29}\text{N}_3\text{O}_5$: 463.2107; found: 463.2119. Calcd. for $\text{C}_{26}\text{H}_{29}\text{N}_3\text{O}_5 \cdot 2.0 \text{ H}_2\text{O}$: C, 62.52; H, 6.65; N, 8.41; found: C, 62.40; H, 6.75, N, 8.62.

Alternatively, similar yields of the title compound can be realized by reaction in refluxing acetone containing a catalytic amount of sodium iodide.

9-Diethylaminoethoxy-20(RS)-camptothecin

15 Hydrochloride. The free base aminoether prepared as described above (275 mg) was suspended in methanol (4 ml) at room temperature and treated dropwise with 0.1N aqueous hydrochloric acid until pH 3 was achieved. The hazy orange solution was evaporated in a nitrogen stream, 20 redissolved in deionized water (50 ml) and filtered through a 0.45 μm membrane. The clear yellow solution was frozen and lyophilized to afford the title compound as a bright yellow fluffy solid (273 mg), mp 232-235°C. γ_{max} (KBr) 2400-3650 (br irregular, OH, amine HCl salt, 25 CH), 1745 (lactone), 1658 (pyridone), 1592 and 1620 (acomatic), 1465, 1400, 1370, 1266, 1232, 1193, 1014, 811 and 720 cm^{-1} ; ^1H NMR ($\text{DMSO}-d_6$) δ 0.89 (t, 3, $J=7$ Hz, H-18), 1.33 (t, 6, $J=7$ Hz, $-\text{N}(\text{CH}_2\text{CH}_3)_2$), 1.88 (m, 2, H-19), 3.34 (m, 4, $-\text{N}(\text{CH}_2\text{CH}_3)_2$), 3.63 (fine t, 2, $-\text{NCH}_2\text{CH}_2\text{O}-$), 30 5.53 (s, 2, H=17), 7.22 (m, 1, H-10), 7.33 (s, 1, H-14), 7.79 (m, 2, H-11 and H-12), 9.05 (s, 1, H-7), 10.46 (br s, 1, $\equiv \text{N}^+\text{H}$). Anal. calcd. for $\text{C}_{26}\text{H}_{30}\text{ClN}_3\text{O}_5 \cdot 1.5 \text{ H}_2\text{O}$: C, 59.26; H, 6.31; N, 7.97; Cl, 6.73. Found: C 59.48; H, 6.27; N, 7.81; Cl, 7.06.

35 Synthesis of 9-(4-methylpiperazine) carbonylamino-2-(RS)-camptothecin Hydrochloride

The title compound was prepared from 9-amino-20(RS)-

camptothecin (synthesis of the amino-camptothecins is described on pp. 33 and 34 of current patent application) in the following manner:

9-(4-Methylpiperazino)carbonylamino-20(RS)-camptothecin.

5 The 9-amino-20(RS)-camptothecin was added to chloroform (treated with alumina to remove hydroxylic components) containing triethylamine. The resulting solution was treated with phosgene gas and filtered to remove solids. The filtrate containing the intermediate
10 carbamoyl chloride was treated with N-methylpiperazine under nitrogen and left overnight. The turbid mixture was washed several times with aqueous sodium bicarbonate solution, dried and evaporated to afford the crude title compound. Chromatography on silica gel provided 9-(4-
15 methylpiperazine)carbonylamino-20(RS)-camptothecin.
9-(4-Methylpiperazino) carbonylamino-20(RS)-camptothecin Hydrochloride.

 The preceding free base urea was suspended in methanol and treated with one equivalent of dilute
20 aqueous hydrochloric acid. The methanol was evaporated and the aqueous residue filtered through a membrane filter. The sample was lyophilized to provide the title compound.

Synthesis of 9-(N,N-Diethylaminoethoxy) carbonylamino-20(RS)-camptothecin Hydrochloride.

 The title compound was prepared from 9-amino-20(RS)-camptothecin in the following manner:

9-(N,N-Diethylaminoethoxy) carbonylamino-20(RS)-camptothecin.

30 The intermediate 9-carbamoyl chloride was prepared as in the preceding example. The resulting chloroform solution was treated with N,N-diethylaminoethanol under nitrogen. After standing overnight, the mixture was washed with aqueous sodium bicarbonate solution, dried
35 and evaporated to afford the crude carbamate. Purification by silica gel chromatography gave the pyure title carbamate as the free base.

9-(N,N-Diethylaminoethoxy)carbonylamino-20(RS)-
camptothecin Hydrochloride.


The free base from the preceding example was
suspended in methanol and treated with one equivalent of
5 dilute aqueous hydrochloric acid. The methanol was
evaporated and the aqueous solution filtered
(membrane). Lyophilization afforded the water soluble
title carbamate.

Obviously, numerous modifications and variations of
10 the present invention are possible in light of the above
teachings. It is therefore to be understood that within
the scope of the appended claims, the invention may be
practiced otherwise than as specifically described
herein.

Claims

1. A method for synthesizing camptothecin and analogs thereof, comprising the steps of:


cyclizing a compound of the formula shown below,
 5 wherein X is an organic group which is converted to a carbonyl group when treated with an acid; in the presence of a basic catalyst, and a source of oxygen,



to form a lactone having the formula



deprotecting said lactone to form a hydroxyl-containing tricyclic compound having the formula shown below, and



15 reacting said hydroxyl-containing tricyclic compound in the presence of an acid or base catalyst, with a substituted ortho-amino compound of the formula



wherein n = 1-2 and wherein each R is selected from
 20 the group consisting of cyano, methylenedioxy, formyl, hydroxy, C₁₋₈ alkoxy, nitro, amino, chloro, bromo, iodo,

fluoro, C₁₋₈ alkyl, trifluoromethyl, aminomethyl, amido, azido and hydrazino groups; R² is H or C₁₋₈ alkyl; and R³ is the side-chain of any of the twenty naturally occurring amino acids.

5 2. The method of Claim 1, wherein said cyclizing step is performed in the presence of a basic catalyst selected from the group consisting of sodium carbonate, potassium carbonate, triethylamine and diisopropylamine.

10 3. The method of Claim 1, wherein said cyclizing step is performed in the presence of a polar organic solvent selected from the group consisting of methanol, ethanol and dimethylformamide.

15 4. The method of Claim 1, wherein said reacting step is catalyzed by a basic compound selected from the group consisting of sodium carbonate, potassium carbonate, triethylamine and diisopropylamine.

20 5. The method of Claim 1, wherein said source of oxygen is selected from the group consisting of oxygen gas, air, hydrogen peroxide, lead tetraacetate and selenium dioxide.

6. The method of Claim 1, wherein R² is H, CH₃ or C₂H₅.

25 7. The method of Claim 1, wherein said reacting step is catalyzed by an acid selected from the group consisting of arylsulfonic acids, acetic acid, formic acid and hydrochloric acid.

30 8. The method of Claim 1, wherein R_n is an amino group, and wherein the amino-camptothecin is further reacted with the carboxylic acid group of an amino acid or carboxylic acid derivative thereof.

9. The method of Claim 1, wherein R_n is an amino group, and wherein the amino-camptothecin is reacted with a C₄₋₁₀ saturated or unsaturated carboxylic acid anhydride.

35 10. The method of Claim 1, wherein R_n is an amino

group at C-9 and wherein the 9-amino-camptothecin is reacted with phosgene and then reacted with a diamine.

11. The method of Claim 10, wherein said diamine is a n-alkylpiperazine.

12. The method of Claim 1, wherein R_n is an amino group at C-9 and wherein the 9-amino-camptothecin is reacted with phosgene and then reacted with a tertiary-amino alcohol.

13. The method of Claim 12, wherein said alcohol is a n-dialkylamino ethanol.

14. The method of Claim 1, wherein R_n is a hydroxy group at C-9 and wherein the 9-hydroxy-camptothecin is reacted with a dialkylamino alkyl halide.


15. The method of Claim 1, wherein R_n is an aminomethyl group and wherein the aminomethyl-camptothecin is reacted with an acid to form an acid addition salt.

16. The method of Claim 15, wherein said acid addition salt is a hydrochloride, gluconate or phosphate.

17. The method of Claim 1, wherein R_n is a hydrazino group and wherein the hydrazino-camptothecin is reacted with an acid to form an acid addition salt.

18. The process of Claim 17, wherein said acid addition salt is a hydrochloride, gluconate or phosphate.

19. A camptothecin analog having the structure shown below



wherein R_n is an amino acid amido group, a C_{4-10} carboxylic acid amido group, a urea group, a urethane group or a dialkylamino ether group or pharmaceutically

acceptable salts thereof.

20. The camptothecin analog of Claim 19, wherein said amino acid amido group is the glycinamido group or pharmaceutically acceptable salts thereof.

5 21. The camptothecin analog of Claim 19, wherein said carboxylic acid amido group is the succinamido group or pharmaceutically acceptable salts thereof.

22. The camptothecin analog of Claim 19, wherein R_n is a urea group or pharmaceutically acceptable salts
10 thereof.

23. The camptothecin analog of Claim 22, wherein said urea group is the N-methylpiperazinocarbonylamino group.

24. The camptothecin analog of Claim 19, wherein R_n
15 is a urethane group or pharmaceutically acceptable salts thereof.

25. The camptothecin analog of Claim 24, wherein the urethane group is the N,N-diethylaminoethoxycarbonylamino group.

20 26. The camptothecin analog of Claim 19, wherein R_n is a dialkylamino ether group.

27. The camptothecin analog of Claim 26, wherein said dialkylamino ether group is the diethylaminoethoxy group.

1/3




FIG. 1A (α)

SUBSTITUTE SHEET

2/3




FIG. 1A(b)

3/3



- a) DMF, K_2CO_3 , 45° , 18h
- b)  45° , 72h
- c) CONCD HCl, GLACIAL HOAc, REFLUX, 2h
- d) ETHYLENE GLYCOL, CH_2Cl_2 , Me_3SiCl , RT, 65h
- e) $(EtO)_2CO$, EtOH, KH, OMe, REFLUX, 3h
- f) $KOtBu$, EtI, DME, $-78^\circ C \longrightarrow$ RT, 18h
- g) Ac_2O , GLACIAL HOAc, H_2 (50psi), RANEY NICKEL, $45^\circ C$, 6h
- h) Ac_2O , HOAc, $NaNO_2$, 0° , 2h
- i) CCl_4 , REFLUX, 8h
- j) K_2CO_3 , MeOH, O_2 , RT, 24h
- k) 2N H_2SO_4 , DME, REFLUX, 24h

FIG. 1B

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US89/04176

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC(5); A61K 31/47; C07D 491/22		
U.S.Cl.: 544/361; 546/48		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U.S.	544/361; 546/48; 514/283	
IPC(5):	A61K 31/47; C07D 491/22	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
Chemical Abstracts CAS ONLINE Search System USPTO APS Search System		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	US, A, 4,473,692 (MIYASAKA, ET AL) 25 September 1984, see the entire document.	19-27
X	US, A, 4,604,463 (MIYASAKA, ET AL) 05 August 1986, see the entire document.	19
A, F	US, A, 4,778,891 (TAGAWA, ET AL) 18 October 1988, see entire document.	1-18
X, E	US, A, 4,894,456 (WALL, ET AL) 16 January 1990, see entire document.	1-27
X	JP, A, 59-051,289 (YAKULT HONSHA CO., LTD.) 24 March 1984, see the entire document.	19
X	JP, A, 59-051,287 (YAKULT HONSHA CO., LTD.) 24 March 1984, see entire document.	19
X	JP, A, 61-050,985 (YAKULT HONSHA CO., LTD.) 13 March 1986, see entire document.	19
X	JP, A, 61-085,319 (YAKULT HONSHA CO., LTD.) 30 April 1986, see entire document.	19
<p>⁹ Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Δ" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
14 FEBRUARY 1990	28 FEB 1990	
International Searching Authority	Signature of Authorized Officer	
ISA/US	 DIANA G. RIVERS	

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

A	EP, A, 220,601 (TAGAWA, ET AL) 06 May 1987, see entire document.	1-18
A	Journal of the American Chemical Society, vol. 94, no. 24, 29 November 1972, Plattner, et al., "Synthesis of Some DE and CDE Ring Analogs of Camptothecin", pages 8613-8615.	1-18
A	Journal of Organic Chemistry, vol. 39, no. 3, 1974, Plattner et al., "Synthesis of Some DE and CDE Ring Analogs of Camptothecin", pages 303-311.	1-18

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE:

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING:

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
X	Journal of Organic Chemistry, vol. 39, no. 23, 1974, Danishefsky, et al., "Synthesis and Biological Evaluation of De- <u>AB</u> -camptothecin" pages 3430-3432.	1-18
X	Journal of Medicinal Chemistry, vol. 23, no. 5, May 1980, Wani, et al., "Plant Antitumor Agents, 18. "Synthesis and Biological Activity of Camptothecin Analogues", pages 554-560.	19, 26-27
X	Chemical Abstracts, vol. 100, no. 17, 1984, Columbus, Ohio, U.S.A., Yang, et al., "Mass Spectrometric Study of Camptothecin and Related Compounds", Abstract no. 139434w, Huaxue Xuebao 1984, 42(1), pp 42-50 (Chinese)	19
X	Journal of Medicinal Chemistry, vol. 29, no. 8, August 1986, Wall, et al., "Plant Antitumor Agents 22. Isolation of 11-Hydroxycamptothecin from Camptotheca acuminata Decne: Total Synthesis and Biological Activity", pages 1553-1555.	1-18
X	Journal of Medicinal Chemistry, vol. 29, no. 11, Nov. 1986, Wani, et al., "Plant Antitumor Agents. 23. Synthesis and Antileukemic Activity of Camptothecin Analogues", pages 2358-2363.	1-19, 26-27